

Pharmacokinetics of oral vitamin C

STEPHEN HICKEY PhD^{i,ii}, HILARY J. ROBERTS PhDⁱⁱⁱ, and NICHOLAS J. MILLER PhD, FRCPath.^{iv}

ⁱ Faculty of Computing, Engineering and Technology, Staffordshire University, Staffordshire, England, ST18 0DG.

ⁱⁱ School of Biology, Chemistry and Health Science, Manchester Metropolitan University, Manchester, England, M1 5GD.

ⁱⁱⁱ Freelance scientific writer.

^{iv} Biolab Medical Unit, 9 Weymouth Street, London, England, W1W 6DB.

Dr Stephen Hickey,
Faculty of Computing, Engineering and Technology,
Staffordshire University,
Staffordshire,
England,
ST18 0DG.
Tel: 01785 353568
Email: s.hickey@staffs.ac.uk

NB: Dr Stephen Hickey is also affiliated with the School of Biology, Chemistry and Health Science, Manchester Metropolitan University, Manchester, England, M1 5GD.

Dr Hilary Roberts
Freelance (unaffiliated)
Contact via Dr Stephen Hickey (see above)

Dr Nicholas J. Miller FRCPath
Biolab Medical Unit
9 Weymouth Street
London W1W 6DB
Tel: 020-7636-5959 / 5905
Fax: 020-7580-3910
E-mail: n.j.miller@biolab.co.uk

Pharmacokinetics of oral vitamin C

STEPHEN HICKEY PhD^{i,ii}, HILARY J. ROBERTS PhDⁱⁱⁱ, and NICHOLAS J. MILLER PhD, FRCPath.^{iv}

ⁱ Faculty of Computing, Engineering and Technology, Staffordshire University, Staffordshire, England, ST18 0DG.

ⁱⁱ School of Biology, Chemistry and Health Science, Manchester Metropolitan University, Manchester, England, M1 5GD.

ⁱⁱⁱ Freelance scientific writer.

^{iv} Biolab Medical Unit, 9 Weymouth Street, London, England, W1W 6DB.

Abstract

Purpose. To test whether plasma vitamin C levels, following oral doses in supplemented volunteers, are tightly controlled and subject to a maximum in the region of 220 $\mu\text{M/L}$, as suggested by previous researchers for depleted subjects.

To determine plasma levels following single, variable-sized doses of standard and liposomal formulations of vitamin C, and compare the effects of the different formulations.

To determine whether plasma levels above about 280 $\mu\text{M/L}$, which have selectively killed cancer, bacteria, or viruses (in laboratory experiments), can be achieved using oral doses of vitamin C.

Design. This was a single blind study, measuring plasma levels in two subjects, in samples taken half-hourly or hourly for six hours, following ingestion of vitamin C. Data were compared with published results and with data from 10 years of laboratory plasma determinations.

Materials and methods. Standard one gram tablets of vitamin C; liposomal vitamin C. Plasma levels were analysed using the method of Butts and Mulvihill.

Results. Preliminary investigations of the effects of liposomal and standard formulation ascorbate showed that blood plasma levels in excess of the previously assumed maximum of 220 $\mu\text{M/L}$ are possible. Large oral doses of liposomal ascorbate resulted in plasma levels above 400 $\mu\text{M/L}$.

Conclusions: Since a single oral dose can produce plasma levels in excess of 400 $\mu\text{M/L}$, pharmacokinetic theory suggests that repeated doses could sustain levels well above the formerly assumed maximum. These results have implications for the use of ascorbate, as a nutrient and as a drug. For example, a short *in vitro* treatment of human Burkitt's lymphoma cells with ascorbate, at 400 $\mu\text{M/L}$, has been shown to result in approximately 50% cancer cell death. Using frequent oral doses, an equivalent plasma level could be sustained indefinitely. Thus, oral vitamin C has potential for use as a non-toxic, sustainable, therapeutic agent.

Further research into the experimental and therapeutic aspects of high, frequent, oral doses of ascorbic acid either alone or (for cancer therapy) in combination with synergistic substances, such as alpha-lipoic acid, copper, or vitamin K3, is needed urgently.

Keywords: Vitamin C, ascorbic acid, nutrition, liposomes, cancer, pharmacokinetics

Correspondence: Dr Steve Hickey, Faculty of Computing, Engineering and Technology,

Staffordshire University, Staffordshire, UK, ST18 0DG.

Email: s.hickey@staffs.ac.uk

Introduction

The amount of vitamin C required for good health represents a challenging aspect of scientific nutrition. Although authorities agree that a few milligrams of ascorbate will prevent acute scurvy, estimates of the optimal intake vary widely. Various experts have proposed doses, ranging from around forty milligrams[1] to several grams a day[2]: a difference more than 100-fold. The debate has been vigorous, especially since Linus Pauling popularised the hypothesis of higher dose requirements.

In an attempt to set vitamin C requirements on a more rigorous scientific basis, the National Institutes of Health (NIH) conducted a series of pharmacokinetic depletion and repletion experiments. These studies were based on seven male[3] and 15 female[4] young, adult subjects.[5] Results indicated that the kidney selectively retains a minimum plasma level of approximately 70 $\mu\text{M/L}$; which varies slightly among subjects. This baseline is protected by ascorbate re-uptake pumps in the kidney, leading to an excretion half-life of 8-40 days, for low (sub-baseline) levels of vitamin C.[6]

Single, daily 200mg doses of vitamin C were shown to maintain this baseline plasma concentration at what the NIH described as a maximal (or “saturation”) level. [4],[5] The NIH reported that single doses higher than about 200mg did not substantially increase the baseline level, which appeared to be limited by the capacity of the re-uptake transporters. The NIH data shows that single, higher, oral doses (up to 1250mg) raise plasma levels transiently.

The shape of the vitamin C response curve results primarily from the rate of oral absorption, combined with an excretion half-life of approximately 30 minutes. A secondary effect concerns its absorption and release from body tissue compartments. Oral absorption of vitamin C is a two-phase, non-linear process; bodily levels are dependent on current intakes. Low intakes produce plasma levels below about 70 $\mu\text{M/L}$, with a long-half life, consistent with resistance to acute scurvy.[7] Higher single doses produce transient increases in plasma concentration, which the NIH group predicted to be limited to a maximum of 220 $\mu\text{M/L}$. [8]

Continuous supplementation, in combination with specific administration technologies, can produce levels greater than the NIH prediction. Here, we provide evidence that plasma levels following oral administration of liposomal ascorbate can reach approximately twice the predicted maximum. These findings have previously unrealised implications for the use of oral vitamin C as a therapeutic agent for various diseases, including cancer.

Methods

Ethical approval for the study was obtained from the ethics committee at Manchester Metropolitan University and the UK Department of Health. Two subjects, one female (52) and one male (55), had previously taken a minimum of six grams per day of vitamin C and an average exceeding 10 grams per day, in divided doses, for a period of 12 months. Both subjects had apparently normal renal function, with no history of kidney stones or renal disease. At the start of the study, each subject fasted for 12 hours,

with no intake of vitamin C before the first (background) measurement. This interval was chosen to ensure almost complete return of the plasma level to baseline, consistent with published pharmacokinetic data,[4],[5],[8] and known absorption and excretion characteristics.[9]

On three sequential mornings, a nurse drew a background blood sample from each subject. Immediately after this, the subjects consumed single 5, 20, or 36 gram doses of vitamin C, in a liposomally encapsulated formulation. Control measurements employed 5 gram doses of a standard ascorbic acid tablet (Table 1). The dosing procedure was single blind: neither nurses nor biochemists were aware of the dose taken. Subsequently, blood samples were taken at half-hourly or hourly intervals by individual venipunctures, to avoid the use of anticoagulants.

[Insert Table 1 about here]

The standard commercial preparation of vitamin C employed nominally one gram tablets with bioflavonoids (Holland and Barrett, Nuneaton, UK). This provides a comparison with existing formulations, since the short-term plasma response to oral ascorbic acid has been quantified previously.[4],[5],[10] The response to these doses was compared with published pharmacokinetic responses to oral doses of ascorbic acid. In addition, one author (NM) had access to the results of plasma ascorbate measurements from Biolab Laboratories. Biolab has made plasma measurements since

1984, and NM has made approximately 30 plasma determinations per week (1500 per year) since 1997; overall about 15,000 measurements.

The liposomal formulation employed nominally one-gram doses, encapsulated in phosphatidylcholine liposomes (Livon Laboratories, Henderson, NV, USA). The liposomal vitamin C was subject to an external high performance liquid chromatography (HPLC: Varian ProStar PDA) validation, which measured 1055mg of ascorbate per (1g) sachet. The measured pH was 6, consistent with the fact that the liposomes contained vitamin C in a salt form, sodium ascorbate.

Blood plasma measurements were obtained over the first six hours of absorption. Plasma vitamin C was measured by the method of Butts and Mulvihill,[11] adapted for the Cobas Mira analyser (F. Hoffman-La Roche Ltd., Diagnostics Division, CH – 4002, Switzerland). The technique depends on reduction of ferric to ferrous ions by ascorbate, which is simultaneously converted to dehydroascorbate. This method had been calibrated against the 2,4-DNPH method[12] and the fluorimetric method.[12] Calibration standards were freshly prepared for each analytical run and three different control samples were included with each batch of samples, giving a between-batch coefficient of variation of approximately 3%.

We analysed the data using standard pattern recognition techniques[13] in terms of outlier determination,[14],[15] and computed the probability that the z value of an

observed pattern was consistent with a normal distribution of values by direct numerical integration, using Simpson's method.[16]

Results

The Biolab plasma ascorbate measurements are entirely consistent with the reference interval of 34 -114 $\mu\text{M/L}$,[17] which corresponds to the 95% interfractile interval of the population studied. Approximately 2.5% of values are below that reference interval, and 2.5% are above the reference interval. Biolab's lowest recorded value for serum vitamin C was 8.0 $\mu\text{M/L}$, at which level the subject was in danger of entering acute scurvy. Of the 15,000 measurements, the highest recorded serum value in a subject not receiving intravenous vitamin C was 220 $\mu\text{M/L}$. This is consistent with the published maximum value. With intravenous administration, Biolab have recorded plasma values in excess of 5000 $\mu\text{M/L}$, in subjects under ascorbate treatment for osteosarcoma.

Baseline levels in the two long-term supplemented subjects, measured at the start of each day, ranged from 100 to 150 $\mu\text{M/L}$. The subjects were thus consistently in the short elimination half-life phase of plasma response.

Figure 1 shows the response of the female subject to single 5g doses of liposomal and standard formulation vitamin C; both produced similar response curves. These results are comparable in form and magnitude to those expected for oral vitamin C in previously depleted subjects. However, peak values exceeded 220 $\mu\text{M/L}$, which has

been reported as the maximum value attainable with repeated oral doses of 3g six times daily.[8] The subjects were experienced users of high-dose vitamin C and neither suffered any gastrointestinal effects at this dose level.

[Insert Figure 1 about here]

Increasing the dose of liposomal vitamin C to 20g gave a broader response, with a delayed maximum, as shown in Figure 2. In this graph, the 20g liposomal dose is compared with a 5g standard dose (male subject). With a 20g intake, the peak plasma level was delayed and the response was broader, indicating a greater absorption of vitamin C. The 5g data set shows a marked outlier (peak): this is attributed to the fact that one of the (5g) blood samples was difficult to extract, with inflammation at the puncture site, providing only a small sample. The subject experienced no bowel tolerance effects at either of these intakes.

[Insert Figure 2 about here]

Figure 3 shows plasma levels following a 36g dose of liposomal vitamin C, for both subjects. This resulted in peak plasma levels, in the region of 400 $\mu\text{M/L}$. A 95% interfractile range (34-114), which contains 95% of the distribution with a mean of 74 corresponds to a calculated standard deviation of 17.4. We note that, under these conditions, an outlier measurement of 400 $\mu\text{M/L}$ would correspond to a deviation of 10.3σ with a theoretical p value of 1.6×10^{-13} (i.e. $P < 0.0000000000001$). With this high

dose, both subjects exceeded their bowel tolerance, leading to diarrhoea. This intolerance presumably arose from the high intake of phospholipid, without food buffering, in fasting individuals. However, our observations using hourly doses suggest that daily intakes of this magnitude are tolerable without bowel effects, as long as the dose is spread throughout the day.

[Insert Figure 3 about here]

Discussion

This is the first published report of the effects of a liposomal formulation on the absorption of vitamin C. This preliminary study covers two subjects, which is approximately 10% of the number used by the NIH to recommend a dietary allowance for the general population. While results from two subjects do not provide an indication of the underlying biodiversity, they demonstrate that previous expectations for vitamin C plasma levels may have been based on a restricted sample of depleted individuals, and do not likely cover the range of responses to oral vitamin C formulations.

It is apparent that the full range of responses in the human population was not represented by the NIH study, as the current data are well outside their published maxima for depleted subjects. These values also exceed statistical expectations,[17] derived from 15,000 subjects measured at Biolab in London. The estimated probability is too small ($P=1.6 \times 10^{-13}$) to be considered accurate but provides a quantitative

indication of the observed discrepancy. Additional research, to include populations not represented in the published data on depleted subjects, is clearly needed.

At over 100 $\mu\text{M/L}$, the subjects' baseline vitamin C levels, following a 12-hour washout period, were higher than those reported in the NIH studies used for the RDA. [4],[5] A possible explanation is that high vitamin C levels redistribute from the blood plasma into other body compartments, and *vice versa*. Thus, subjects who habitually consume large amounts of vitamin C might carry a reserve in their tissues. Plasma vitamin C, which is depleted through kidney excretion, has a half-life of only half an hour. However, while it is being excreted, it can be replaced by absorption from the gut, or by redistribution from other body compartments.

The elevated baselines observed in these subjects are inconsistent with the NIH's pharmacokinetic modelling of the plasma response of gram level doses of vitamin C.[3], [4],[5] Regular use of high dose vitamin C, given at intervals of 12 hours or less, produces baseline plasma levels greater than those the NIH study described as "saturated". Since "saturation" implies a maximum level, which has been refuted by the higher levels reported here, use of this term (as employed by the NIH in the context of vitamin C plasma levels) should be considered obsolete, inappropriate, and inaccurate, and the word returned to its conventional meaning.

The plasma levels reported here are higher than is usually seen with oral administration of vitamin C, although the doses were also higher. Biolab noted that the

plasma levels attained with 20g and 36g doses of liposomal vitamin C were higher than any they had measured previously, following oral doses. Biolab's previous experience and measurement of plasma vitamin C levels has been consistent with the reference interval for human plasma (34-114 $\mu\text{M/L}$). Neonates and supplemented subjects with impaired renal function routinely record plasma ascorbate concentrations of up to 200 $\mu\text{M/L}$.

Since plasma levels in this study were outside the normally reported range, they were communicated to Dr Ron Hunninghake, Clinical Director of the International Center for the Improvement Of Human Functioning, who carried out an independent replication. Hunninghake, who acted as his own experimental subject, is a habitual high-dose vitamin C supplement user (10 grams per day). Using repeated doses of liposomal and standard vitamin C, at lower intakes than our 36g maximum, Hunninghake determined vitamin C levels in excess of 300 $\mu\text{M/L}$. This level was achieved with 10g of standard C, followed by an additional 12 or 18 grams of liposomal C, divided at intervals through the day. Through this procedure, high vitamin C plasma levels were sustained, rather than peaking as in our single dose study. Following a washout period (no vitamin C intake), Hunninghake found he retained between 100-150 $\mu\text{M/L}$, which supports our findings.

Hunninghake's laboratory (Bio-Center, Wichita, USA) has averaged approximately 2000 plasma C determinations each year since 1994; in 2006, they performed 6634 plasma measurements. Their background results (from 37-120 $\mu\text{M/L}$)

provide a reference range that is consistent with Biolab's (34 -114 $\mu\text{M/L}$). The maximum value recorded at the Bio-Centre (from a cancer patient, taking approximately 100g per day orally) was 318 $\mu\text{M/L}$. [18]

The results of the present study, together with Hunninghake's replication, clearly refute the NIH group's suggestion that the blood is saturated with vitamin C at around 70 $\mu\text{M/L}$.

The similarity in the plasma response curves for liposomal vitamin C and the standard commercial formulation, shown in Figure 1 (5g doses), is interesting. There is a hint that the liposomal form has a slower onset to peak level and a broader profile. Liposomes are absorbed from the gut and into the liver, before being released into the blood stream. This response can be seen more clearly in the 20g dose, in Figure 2. It is apparent that sustained levels of plasma ascorbate, above the previously assumed maximum of 220 $\mu\text{M/L}$, are possible with oral intakes of liposomal vitamin C.

Cathcart describes a mechanism of increased bowel tolerance to oral ascorbate. [19] He reports a direct correlation between the severity of illness and the maximum amount of ascorbate absorbed by the oral route. These clinical observations indicate that the physiological response to ascorbate is variable; clarification awaits suitable pharmacokinetic studies in diseased patients.

The results of this preliminary study are consistent with the suggestion that continuous vitamin C supplementation facilitates increased absorption of large doses. The reference interval for vitamin C quoted for human plasma is 34 to 114 $\mu\text{M/L}$. [17] In these experiments, the measured baseline levels for vitamin C were in the range 132 to 149 $\mu\text{M/L}$. This concentration is approximately twice the level found in previously depleted subjects. The mechanisms of absorption of ascorbate from the gut are not well understood. Our experience suggests that subjects who habitually consume high intakes of oral vitamin C and subjects that are ill may show a differing response to those who were previously depleted. Physiological adaptation to ascorbate intake may involve recruitment of glucose transporters (GLUT4) signalled by insulin, cellular contraction, or reactive oxygen species. [20],[21]

Our observations suggest that poor oral absorption is related to carbohydrate intake. Glucose and other sugars compete with dehydroascorbate for absorption by GLUT transporters. The subjects in this study reported that they often consumed 10-grams, or more, of ascorbate in a single dose, without significant bowel effects, provided their carbohydrate intake was low. In the present study, the subjects reported no bowel effects, or a slight constipation, following the 5-gram doses. The 20-gram liposomal dose was tolerated well. In these fasting individuals, the bowel reaction to the 36-gram dose may have been caused by the large intake of phospholipids, without food buffering.

The pharmacokinetics that form a principal evidence base for determination of the RDA values for vitamin C rely on results from a small number of depleted, or previously depleted, individuals. Such individuals may not be representative of all members of the population. Well-nourished individuals, with western levels of carbohydrate intake, who have been previously depleted of ascorbate, may have a low plasma response to oral ascorbate. In the present study, the levels attained from a single oral dose of five standard one-gram tablets produced plasma levels above the predicted 220 $\mu\text{M/L}$ maximum level. Larger or repeated doses would have increased this level further.

The “saturated” baseline level, determined as 70-80 $\mu\text{M/L}$ in the NIH pharmacokinetic studies, is not an upper limit. Baseline levels between 12 and 24 hours after a single oral dose are consistent with ascorbate transfer between blood plasma and other body compartments. In our study, the recorded background level after a 12-hour washout was in the range 132 - 149 $\mu\text{M/L}$, approximately double that found in depleted individuals. This is consistent with increased tissue levels or sustained absorption from the intestine. We suggest the NIH’s assumption that plasma levels are tightly controlled and “saturated” at 70-80 $\mu\text{M/L}$ was premature, and thus the RDA for ascorbate could require revision.

Our findings have important implications for therapy. The role of pharmacological levels of nutrients in cancer and other diseases is currently being re-evaluated.[22] In particular, the cytotoxic action of ascorbate on cancer cells suggests

the possibility of a potentially non-toxic cancer therapy.[23],[24] However, it has been assumed that cytotoxic levels of ascorbate could only be achieved with intravenous infusions of sodium ascorbate.[8] This would be an implication of the available data, if oral doses could achieve only 220 $\mu\text{M/L}$ in plasma. However, we have demonstrated that single doses of liposomal formulations can give levels above 400 $\mu\text{M/L}$. If given in a single dose to a fasting individual, such intakes might be impractical. However, these preliminary findings suggest that plasma levels of 500-600 $\mu\text{M/L}$, or more, could be sustained indefinitely with smaller, but repeated, oral intakes.

Data for the cytotoxicity of ascorbate comes from laboratory experiments, performed over a timescale measured in hours. However, tumours can selectively concentrate ascorbate from plasma. Published results for a one-hour *in vitro* treatment of human Burkitt's lymphoma cells, using ascorbate at 300 $\mu\text{M/L}$, show approximately 30% necrosis and apoptosis; 400 $\mu\text{M/L}$ increases the cell death to approximately 50%. [25] Moreover, ascorbate cytotoxicity is synergistic with other oral nutrients, including alpha-lipoic acid,[26] vitamin K3,[27] and copper.[28] The implications of sustained high concentrations of ascorbate may thus be therapeutically significant.

References

- [1] Dietary Reference Values for Food Energy and Nutrients for the United Kingdom., 1991, Report of the Panel on Dietary Reference Values, Committee on Medical Aspects of Food and Nutrition Policy, HMSO, London.
- [2] Pauling L. 2006, How to Live Longer And Feel Better, Corvallis: Oregon State University Press.
- [3] Levine M. Conry-Cantilena C. Wang Y. Welch R.W. Washko P.W. Dhariwal K.R. Park J.B. Lazarev A. Graumlich J.F. King J. Cantilena L.R., 1996, Vitamin C pharmacokinetics in healthy volunteers: evidence for a recommended dietary allowance, *Proc Natl Acad Sci USA*, **93**(8), 3704-9.
- [4] Levine M. Wang Y. Padayatty S.J. Morrow J., 2001, A new recommended dietary allowance of vitamin C for healthy young women, *Proc Natl Acad Sci USA*, **98**(17):9842-6.
- [5] Levine M. Rumsey S.C. Daruwala R. Park J.B. Wang Y., 1999, Criteria and recommendations for vitamin C intake, *JAMA*, **281**(15), 1415-23.
- [6] Kallner A. Hartmann D. Hornig D., 1979, Steady-state turnover and body pool of ascorbic acid in man, *Am J Clin Nutr*, **32**, 530-539.
- [7] Hickey S. Roberts H.J. Cathcart R.F., 2005, Dynamic flow, *JOM*, **20**(4), 237-244.
- [8] Padayatty S.J. Sun H. Wang Y. Riordan H.D. Hewitt S.M. Katz A. Wesley R.A. Levine M., 2004, Vitamin C Pharmacokinetics: Implications for Oral and Intravenous Use, *Ann Intern Med*, **140**, 533-537.
- [9] Hickey S. Roberts H., 2004, *Ascorbate: The Science of Vitamin C*, Lulu press.
- [10] Benke K.K., 1999, Modelling Ascorbic Acid Level in Plasma and Its Dependence on Absorbed Dose, *Journal of the Australasian College of Nutritional & Environmental Medicine*, **18**(1), 11-12.
- [11] Butts W.C. Mulvihill H.J., 1975, Centrifugal analyzer determination of ascorbate in serum or urine with Fe³⁺/ferrozine. *Clin Chem*, **21**, 493-1487.
- [12] Deutsch M.J. Weeks C.E., 1965, Microfluorimetric assay for vitamin C. *J of the Assoc. of Official Analytical Chemists*, **48**, 1248-1256.
- [13] Krishnaiah P.R. Kanal L.N., 1982, Classification, Pattern Recognition and Reduction of Dimensionality, *Handbook of Statistics*, Vol 2, North Holland, New York.
- [14] Duda R.O. Hart P.E., 1973, *Pattern Classification and Scene Analysis*, John Wiley & Sons, New York.

-
- [15] Sæbø H.V. Bråten K. Hjort N.L. Llewellyn B. Moen E., 1985, Contextual classification of remotely sensed data: Statistical methods and development of a system, Report no. 768, Norwegian Computing Center.
- [16] Press W.H. Flannery B.P. Teukolsky S.A. Vetterling W.T., 1992, Numerical Recipes in C: The Art of Scientific Computing, Cambridge University Press.
- [17] Brewster M.A. Turley C.P., 1987, Vitamin C. In: Pesce A.J. and Kaplan L.A., eds. Methods in Clinical Chemistry. St. Louis: C.V. Mosby Co., 574-581.
- [18] Hunninghake R., 2008, Conquer Cancer with Redox Synergy, 37th Annual Conference International Society for Orthomolecular Medicine, Vancouver, May, (personal communication before presentation).
- [19] Cathcart R.F. Vitamin C, titrating to bowel tolerance, anascorbemia, and acute induced scurvy, Med Hypotheses, **7**(11), 1359-1376.
- [20] Lund S. Holman G.D., 1995, Schmitz O. Pedersen O. Contraction stimulates translocation of glucose transporter GLUT4 in skeletal muscle through a mechanism distinct from that of insulin, Proc Natl Acad Sci USA, **92**(13), 5817-21.
- [21] Sandström M.E. Zhang S.J. Bruton J. Silva J.P. Reid M.B. Westerblad H. Katz A., 2006, Role of reactive oxygen species in contraction-mediated glucose transport in mouse skeletal muscle, J Physiol, **575**(1), 251-262.
- [22] Padayatty S.J. Riordan H.D. Hewitt S.M. Katz A. Hoffer L.J. Levine M., 2006, Intravenously administered vitamin C as cancer therapy: three cases, CMAJ, **174**(7), 937-942.
- [23] Padayatty S.J. Levine M., 2000, Reevaluation of ascorbate in cancer treatment: emerging evidence, open minds and serendipity, J Am Coll Nutr, **19**(4), 423-425.
- [24] Gonzalez M.J. Miranda-Massari J.R. Mora E.M. Guzman A. Riordan N.H. Riordan H.D. Casciari J.J. Jackson J.A., 2005, Roman-Franco A. Orthomolecular oncology review: ascorbic acid and cancer 25 years later, Integr Cancer Ther, **4**(1), 32-44.
- [25] Chen Q. Espey M.G. Krishna M.C. Mitchell J.B. Corpe C.P. Buettner G.R. Shacter E. Levine M., 2005, Pharmacologic ascorbic acid concentrations selectively kill cancer cells: action as a pro-drug to deliver hydrogen peroxide to tissues, Proc Natl Acad Sci USA, **102**(38), 13604-13609.
- [26] Casciari J.J. Riordan N.H. Schmidt T.L. Meng X.L. Jackson J.A. Riordan H.D., 2001, Cytotoxicity of ascorbate, lipoic acid, and other antioxidants in hollow fibre in vitro tumours, Br J Cancer, **84**(11), 1544-1550.

[27] Calderon P.B. Cadrobbi J. Marques C. Hong-Ngoc N. Jamison J.M. Gilloteaux J. Summers J.L. Taper H.S., 2002, Potential therapeutic application of the association of vitamins C and K3 in cancer treatment, *Curr Med Chem*, **9**(24), 2271-2285.

[28] González MJ, Mora EM, Miranda-Massari JR, Matta J, Riordan HD, Riordan NH., 2002, Inhibition of human breast carcinoma cell proliferation by ascorbate and copper, *P R Health Sci J*, **21**(1), 21-23.

Table 1

Protocol for doses			
	Day 1	Day 2	Day 3
Subject 1 (M)	5g (stn)	20g (lip)	36g (lip)
Subject 2 (F)	5g (lip)	5g (stn)	36g (lip)

Figure 1




Figure 2




Figure 3




Figure captions

Figure 1. Plasma response curve following 5g oral doses of standard vitamin C tablets (o) and liposomal ascorbate (Δ) in the same subject.

Figure 2. Plasma response to single doses: 5g standard (Δ) and 20g liposomal (o) ascorbate. (NB: Sampling problem - only a small blood sample was able to be taken at 275 minutes, corresponding to the outlier.)

Figure 3. Plasma response to a single 36g dose of liposomal ascorbate in two subjects.